

One-instanton predictions of Seiberg-Witten curves for product groups

Isabel P. Ennes¹

Martin Fisher School of Physics
Brandeis University, Waltham, MA 02454

Stephen G. Naculich²

Department of Physics
Bowdoin College, Brunswick, ME 04011

Henric Rhedin

Martin Fisher School of Physics
Brandeis University, Waltham, MA 02454
and

Department of Engineering Sciences³,
Physics and Mathematics
Karlstad University, S-651 88 Karlstad, Sweden

Howard J. Schnitzer⁴

Martin Fisher School of Physics³
Brandeis University, Waltham, MA 02454
and

Lyman Laboratory of Physics
Harvard University, Cambridge, MA 02138

Abstract

One-instanton predictions for the prepotential are obtained from the Seiberg-Witten curve for the Coulomb branch of $\mathcal{N} = 2$ supersymmetric gauge theory for the product group $\prod_{n=1}^m \text{SU}(N_n)$ with a massless matter hypermultiplet in the bifundamental representation $(N_n, \bar{N}_{n+1})$ of $\text{SU}(N_n) \times \text{SU}(N_{n+1})$ for $n = 1$ to $m - 1$, together with N_0 and N_{m+1} matter hypermultiplets in the fundamental representations of $\text{SU}(N_1)$ and $\text{SU}(N_m)$ respectively. The derivation uses a generalization of the systematic perturbation expansion about a hyperelliptic curve developed by us in earlier work.

¹Research supported by the DOE under grant DE-FG02-92ER40706.

²Research supported in part by the National Science Foundation under grant no. PHY94-07194.

³Permanent address

⁴Research supported in part by the DOE under grant DE-FG02-92ER40706.

naculich@bowdoin.edu; henric.rhedin@kau.se; ennes,schnitzer@binah.cc.brandeis.edu

Spectacular advances have been made in our understanding of the non-perturbative behavior of supersymmetric gauge theories and string theories. In particular, the program of Seiberg and Witten [1] allows one to compute the exact behavior of low-energy four-dimensional $\mathcal{N} = 2$ supersymmetric gauge theories in various regions of moduli space from the following input: a Riemann surface or algebraic curve appropriate to the specific theory, and the Seiberg-Witten meromorphic one-form. When the curve in question is non-hyperelliptic, however, the explicit extraction of this information is a challenging technical problem. In this letter, we will use Seiberg-Witten theory to calculate the one-instanton predictions of an $\mathcal{N} = 2$ supersymmetric gauge theory based on the product group $\prod_{n=1}^m \text{SU}(N_n)$ by extending the methods of ref. [2]-[7].

The exact low-energy properties of $\mathcal{N} = 2$ theories are encapsulated in the form of the prepotential $\mathcal{F}(A)$, in terms of which the Wilson effective Lagrangian is

$$\mathcal{L} = \frac{1}{4\pi} \text{Im} \left[\int d^4\theta \frac{\partial \mathcal{F}(A)}{\partial A^i} \bar{A}^i + \frac{1}{2} \int d^2\theta \frac{\partial^2 \mathcal{F}(A)}{\partial A^i \partial A^j} W^i W^j \right], \quad (1)$$

to lowest order in the momentum expansion, where A^i are $\mathcal{N} = 1$ chiral superfields and W^i are $\mathcal{N} = 1$ vector superfields. Holomorphy implies that the prepotential in the Coulomb phase has the form of an instanton expansion

$$\mathcal{F}(A) = \mathcal{F}_{\text{cl}}(A) + \mathcal{F}_{1\text{-loop}}(A) + \sum_{d=1}^{\infty} \mathcal{F}_{d\text{-inst}}(A). \quad (2)$$

Consider an $\mathcal{N} = 2$ supersymmetric gauge theory based on the gauge group $\prod_{n=1}^m \text{SU}(N_n)$. In addition to the chiral gauge multiplet in the adjoint representation of each of the factor groups, the theory we are considering contains a massless matter hypermultiplet in the bifundamental representation $(N_n, \bar{N}_{n+1})$ of $\text{SU}(N_n) \times \text{SU}(N_{n+1})$ for $n = 1, \dots, m-1$; N_0 matter hypermultiplets in the fundamental representation of $\text{SU}(N_1)$ (whose masses we denote $e_k^{(0)}, 1 \leq k \leq N_0$); and N_{m+1} matter hypermultiplets in the fundamental representation of $\text{SU}(N_m)$ (whose masses we denote $e_k^{(m+1)}, 1 \leq k \leq N_{m+1}$). The adjoint multiplets contain complex scalar fields $\phi^{(n)}$ for each of the factor groups. The Lagrangian has a potential with flat directions, along which the

symmetry is generically broken to $\prod_{n=1}^m \text{U}(1)^{N_n-1}$. The moduli space of the theory is therefore parametrized by the order parameters $e_k^{(n)}$ ($1 \leq k \leq N_n$, $1 \leq n \leq m$), which are the eigenvalues of the $\phi^{(n)}$ and satisfy the constraint $\sum_{k=1}^{N_n} e_k^{(n)} = 0$.

The curve for this product group theory was obtained using M-theory [8] and geometric engineering [9], and made more explicit in ref. [10]:

$$P_{N_0}(x)t^{m+1} - P_{N_1}(x)t^m + \sum_{j=0}^{m-1} (-)^{m-j+1} \left[\prod_{\ell=1}^{m-j} L_\ell^{m-\ell-j+1} \right] P_{N_{m-j+1}}(x)t^j = 0, \quad (3)$$

where

$$P_{N_n}(x) = \prod_{i=1}^{N_n} (x - e_i^{(n)}), \quad (n = 0 \text{ to } m),$$

$$L_n^2 = \Lambda_n^{2N_n - N_{n-1} - N_{n+1}}, \quad (4)$$

with Λ_n the quantum scale of the gauge group $\text{SU}(N_n)$. The requirement of asymptotic freedom, and restriction to the Coulomb phase, implies that Λ_n appear with positive powers in (3).

The curve (3) describes a $(m+1)$ -fold branched covering of the Riemann sphere, with sheets n and $n+1$ connected by N_n square-root branch-cuts centered about $x = e_k^{(n)}$ ($k = 1$ to N_n), and having endpoints $x_k^{(n)-}$ and $x_k^{(n)+}$. Following the approach of Seiberg-Witten, we will use this curve to compute the renormalized order parameters and their duals

$$2\pi i a_k^{(n)} = \oint_{A_k^{(n)}} \lambda \quad \text{and} \quad 2\pi i a_{D,k}^{(n)} = \oint_{B_k^{(n)}} \lambda, \quad (5)$$

where λ is the Seiberg-Witten differential, and $A_k^{(n)}$ and $B_k^{(n)}$ ($2 \leq k \leq N_n$) are a set of canonical homology cycles for the Riemann surface. The cycle $A_k^{(n)}$ is chosen to be a simple contour on sheet n enclosing the branch cut centered about $e_k^{(n)}$. The cycle $B_k^{(n)}$ goes from $x_1^{(n)-}$ to $x_k^{(n)-}$ on the n^{th} sheet and from $x_k^{(n)-}$ to $x_1^{(n)-}$ on the $(n+1)^{\text{th}}$ [2]. Once we obtain $a_k^{(n)}$ and $a_{D,k}^{(n)}$, the prepotential can be computed by integrating

$$a_{D,k}^{(n)} = \frac{\partial \mathcal{F}}{\partial a_k^{(n)}}, \quad (k = 2 \text{ to } N_n, \quad n = 1 \text{ to } m). \quad (6)$$

In our computation, we will perform a multiple perturbation expansion in the several quantum scales L_n , with the result for the prepotential given to one-instanton accuracy, *i.e.*, $O(L_n^2)$ for all n . To calculate $a_k^{(n)}$ and $a_{D,k}^{(n)}$ for the group $SU(N_n)$, we define $t = \tilde{t} \prod_{\ell=1}^{n-1} L_\ell^2$ to recast the curve (3) as

$$\sum_{j=0}^n (-)^j \left[\prod_{\ell=j+1}^{n-1} L_\ell^{2(\ell-j)} \right] P_{N_j}(x) \tilde{t}^{m+1-j} + \sum_{j=n+1}^{m+1} (-)^j \left[\prod_{\ell=n}^{j-1} L_\ell^{2(j-\ell)} \right] P_{N_j}(x) \tilde{t}^{m+1-j} = 0, \quad (7)$$

or more explicitly

$$\begin{aligned} \dots &+ (-)^{n-3} L_{n-2}^2 L_{n-1}^4 P_{N_{n-3}}(x) \tilde{t}^{m-n+4} + (-)^{n-2} L_{n-1}^2 P_{N_{n-2}}(x) \tilde{t}^{m-n+3} \\ &+ (-)^{n-1} P_{N_{n-1}}(x) \tilde{t}^{m-n+2} + (-)^n P_{N_n}(x) \tilde{t}^{m-n+1} + (-)^{n+1} L_n^2 P_{N_{n+1}}(x) \tilde{t}^{m-n} \\ &+ (-)^{n+2} L_n^4 L_{n+1}^2 P_{N_{n+2}}(x) \tilde{t}^{m-n-1} + (-)^{n+3} L_n^6 L_{n+1}^4 L_{n+2}^2 P_{N_{n+3}}(x) \tilde{t}^{m-n-2} + \dots = 0. \end{aligned} \quad (8)$$

To obtain the one-loop, zero-instanton contribution, *i.e.*, $O(\log L_n)$, to $a_k^{(n)}$ and $a_{D,k}^{(n)}$, one may set $L_\ell = 0$ for $\ell \neq n$, in which case the curve (8) reduces, after the change of variable $\tilde{t} = y/P_{N_{n-1}}(x)$, to the hyperelliptic curve

$$y^2 + 2A(x)y + B(x) = 0, \quad (9)$$

with

$$\begin{aligned} A(x) &= -\frac{1}{2} P_{N_n}(x), \\ B(x) &= L_n^2 P_{N_{n+1}}(x) P_{N_{n-1}}(x). \end{aligned} \quad (10)$$

On one of the sheets, eq. (9) has the solution

$$y = -A - r \quad \text{where} \quad r = \sqrt{A^2 - B}, \quad (11)$$

from which we may compute the Seiberg-Witten differential $\lambda = xdy/y$ in the hyperelliptic approximation to be

$$\lambda_I = \frac{x \left(\frac{A'}{A} - \frac{B'}{2B} \right)}{\sqrt{1 - \frac{B}{A^2}}} dx. \quad (12)$$

(On the other sheet, the solution to eq. (9) is $y = -A + r$).

To obtain the one-instanton correction (*i.e.*, $O(L_\ell^2)$ for all ℓ) to the order parameters, one again makes the change of variables $\tilde{t} = y/P_{N_{n-1}}(x)$ in eq. (8), and keeps two more terms beyond those in eq. (9), obtaining the quartic curve

$$\epsilon_1(x)y^4 + y^3 + 2A(x)y^2 + B(x)y + \epsilon_2(x) = 0, \quad (13)$$

where

$$\begin{aligned} \epsilon_1(x) &= -L_{n-1}^2 \frac{P_{N_{n-2}}(x)}{P_{N_{n-1}}^2(x)}, \\ \epsilon_2(x) &= -L_n^4 L_{n+1}^2 P_{N_{n+2}}(x) P_{N_{n-1}}^2(x). \end{aligned} \quad (14)$$

Rewriting eq. (13) as

$$y + A + r = \frac{1}{y + A - r} \left[-\epsilon_1 y^3 - \frac{\epsilon_2}{y} \right] \quad (15)$$

and substituting $y = -A - r$ into the right hand side, which is already first order in ϵ , we obtain

$$y = -A - r - \frac{(A+r)^3}{2r} \epsilon_1 - \frac{1}{2r(A+r)} \epsilon_2 + \dots, \quad (16)$$

to first order in ϵ_1 and ϵ_2 . The Seiberg-Witten differential is correspondingly modified to

$$\lambda = \lambda_I + \lambda_{II} + \dots, \quad (17)$$

where λ_I is the hyperelliptic approximation (12) and

$$\lambda_{II} = \left(-A \epsilon_1 - \frac{A}{B^2} \epsilon_2 \right) dx = -L_{n-1}^2 \frac{P_{N_n} P_{N_{n-2}}}{2P_{N_{n-1}}^2} dx - L_{n+1}^2 \frac{P_{N_n} P_{N_{n+2}}}{2P_{N_{n+1}}^2} dx, \quad (18)$$

obtained from a calculation similar to that in Appendix C of ref. [5].

One computes the order parameters (5) using the methods of refs. [2, 5, 6], obtaining

$$a_k^{(n)} = e_k^{(n)} + \frac{1}{4} L_n^2 \frac{\partial S_k^{(n)}}{\partial x} (e_k^{(n)}) + \dots, \quad (k = 1 \text{ to } N_n, \quad n = 1 \text{ to } m), \quad (19)$$

where the residue functions $S_k^{(n)}(x)$ are defined in terms of eq. (10) by

$$\frac{S_k^{(n)}(x)}{(x - e_k^{(n)})^2} = \frac{B(x)}{A(x)^2}. \quad (20)$$

Considerations analogous to those of Appendix D of ref. [5] give the identities

$$\sum_{j=1}^{N_n} \frac{\partial S_j^{(n)}}{\partial x}(e_j^{(n)}) = 0, \quad (21)$$

implying $\sum_{i=1}^{N_n} a_i^{(n)} = \sum_{i=1}^{N_n} e_i^{(n)}$ to the order that we are working.

Next, the dual order parameters $a_{D,k}^{(n)}$ are computed along the lines of sec. 5 of ref. [5], giving

$$\begin{aligned} 2\pi i a_{D,k}^{(n)} &= [2N_n - N_{n+1} - N_{n-1} + 2 \log L_n + \text{const}] a_k^{(n)} - 2 \sum_{i \neq k}^{N_n} (a_k^{(n)} - a_i^{(n)}) \log(a_k^{(n)} - a_i^{(n)}) \\ &+ \sum_{i=1}^{N_{n+1}} (a_k^{(n)} - a_i^{(n+1)}) \log(a_k^{(n)} - a_i^{(n+1)}) + \sum_{i=1}^{N_{n-1}} (a_k^{(n)} - a_i^{(n-1)}) \log(a_k^{(n)} - a_i^{(n-1)}) \\ &+ \frac{1}{4} L_n^2 \frac{\partial S_k^{(n)}}{\partial x}(a_k^{(n)}) - \frac{1}{2} L_n^2 \sum_{i \neq k}^{N_n} \frac{S_i^{(n)}(a_i^{(n)})}{a_k^{(n)} - a_i^{(n)}} + \frac{1}{4} L_{n+1}^2 \sum_{i=1}^{N_{n+1}} \frac{S_i^{(n+1)}(a_i^{(n+1)})}{a_k^{(n)} - a_i^{(n+1)}} \\ &+ \frac{1}{4} L_{n-1}^2 \sum_{i=1}^{N_{n-1}} \frac{S_i^{(n-1)}(a_i^{(n-1)})}{a_k^{(n)} - a_i^{(n-1)}} + \dots, \quad (k = 2 \text{ to } N_n, \quad n = 1 \text{ to } m). \end{aligned} \quad (22)$$

In eq. (22), we define $a_k^{(0)} = e_k^{(0)}$ and $a_k^{(m+1)} = e_k^{(m+1)}$ (the masses of the hypermultiplets in the fundamentals of $SU(N_1)$ and $SU(N_m)$ respectively), and $L_0 = L_{m+1} = 0$.

One then integrates eq. (6) using eq. (22) to obtain the prepotential (2) to one-instanton accuracy, finding

$$\begin{aligned} \mathcal{F}_{1\text{-loop}} &= \frac{i}{8\pi} \sum_{n=1}^m \sum_{i,j=1}^{N_n} (a_i^{(n)} - a_j^{(n)})^2 \log(a_i^{(n)} - a_j^{(n)})^2 \\ &- \frac{i}{8\pi} \sum_{n=0}^m \sum_{i=1}^{N_n} \sum_{j=1}^{N_{n+1}} (a_i^{(n)} - a_j^{(n+1)})^2 \log(a_i^{(n)} - a_j^{(n+1)})^2, \end{aligned} \quad (23)$$

and

$$\mathcal{F}_{1\text{-inst}} = \frac{1}{8\pi i} \sum_{n=1}^m L_n^2 \sum_{k=1}^{N_n} S_k^{(n)}(a_k^{(n)}), \quad (24)$$

where

$$S_k^{(n)}(x) = \frac{4 \prod_{i=1}^{N_{n+1}} (x - a_i^{(n+1)}) \prod_{i=1}^{N_{n-1}} (x - a_i^{(n-1)})}{\prod_{i \neq k}^{N_n} (x - a_i^{(n)})^2}, \quad (k = 1 \text{ to } N_n, \quad n = 1 \text{ to } m). \quad (25)$$

Note that as $S_k^{(n)}(a_k^{(n)})$ depends on $a_i^{(n+1)}$ and $a_i^{(n-1)}$ as well as $a_i^{(n)}$, eq. (24) is *not* just the naive sum of instanton contributions from each subgroup.

The one-loop prepotential (23) agrees with the perturbation theory result for a chiral gauge multiplet in the adjoint representation of each of the factor groups, a massless matter hypermultiplet in the bifundamental representation $(N_n, \bar{N}_{n+1})$ of $SU(N_n) \times SU(N_{n+1})$ for $n = 1$ to $m-1$, N_0 matter hypermultiplets with masses $a_k^{(0)}$ in the fundamental representation of $SU(N_1)$, and N_{m+1} matter hypermultiplets with masses $a_k^{(m+1)}$ in the fundamental representation of $SU(N_m)$.

One check of the one-instanton correction (24) is provided by ref. [4], where various decoupling limits for $\mathcal{N} = 2$ $SU(N)$ gauge theory with a massive hypermultiplet in the adjoint representation are considered. D'Hoker and Phong [4] obtain $\mathcal{F}_{1-\text{inst}}$ for the product group theory, but with restriction to a single quantum scale. We find agreement with their result when we restrict eqs. (24) and (25) to the special case of a single quantum scale, which therefore provides a test of the curve (3) obtained from M-theory.

In this paper, we showed that to compute the order parameters of the $\mathcal{N} = 2$ gauge theory for the product group $\prod_{n=1}^m SU(N_n)$ to one-instanton accuracy, one need only consider the sequence of quartic curves (13), even though the complete curve for the theory (3) is of higher order (*viz.*, $m+1$) for $m > 3$, *i.e.*, for products of three or more groups. (The case $m = 2$ was analyzed in ref. [7].) In the language of type IIA string theory, this means one need only consider all possible chains of four parallel neighboring NS 5-branes, among the total set of $m+1$ parallel NS 5-branes, to achieve one-instanton accuracy. For higher instanton accuracy, additional parallel 5-branes are required. An analogue of this result plays a crucial role in our analysis of the prepotential and Seiberg-Witten curve for $SU(N)$ gauge theory with two antisymmetric and N_f fundamental hypermultiplets [11].

Acknowledgement: HJS wishes to thank the Physics Department of Harvard University for their continued hospitality, and to the CERN theory group for hospitality during summer 1998.

References

- [1] N. Seiberg and E. Witten, Nucl. Phys. B426 (1994) 19, hep-th/9407087; B430 (1994) 485(E); Nucl. Phys. B431 (1994) 484, hep-th/9408099.
- [2] E. D'Hoker, I.M. Krichever, and D.H. Phong, Nucl. Phys. B489 (1997) 179, hep-th/9609041.
- [3] E. D'Hoker, I.M. Krichever, and D.H. Phong, Nucl. Phys. B489 (1997) 211, hep-th/9609145; Nucl. Phys. B494 (1997) 89, hep-th/9610156; E. D'Hoker and D.H. Phong, Phys. Lett. B397 (1997) 94, hep-th/9701055.
- [4] E. D'Hoker and D.H. Phong, Nucl. Phys. B513 (1998) 405, hep-th/9709053.
- [5] S.G. Naculich, H. Rhedin, and H.J. Schnitzer, Nucl. Phys. B533 (1998) 275, hep-th/9804105.
- [6] I.P. Ennes, S.G. Naculich, H. Rhedin, and H.J. Schnitzer, hep-th/9804151, Int. Jour. Mod. Phys. A, in press.
- [7] I.P. Ennes, S.G. Naculich, H. Rhedin, and H.J. Schnitzer, Nucl. Phys. B536 (1998) 245, hep-th/9806144.
- [8] E. Witten, Nucl. Phys. B500 (1997) 3, hep-th/9703166.
- [9] S. Katz, P. Mayr, and C. Vafa, Adv. Theor. Math. Phys. 1 (1998) 53, hep-th/9706110.
- [10] J. Erlich, A. Naqvi, and L. Randall, Phys. Rev. D58 (1998) 046002, hep-th/9801108.
- [11] I.P. Ennes, S.G. Naculich, H. Rhedin, and H.J. Schnitzer, BRX-TH-447, BOW-PH-114, HUTP-98/A085, to appear.